

LATENT TUBERCULOSIS INFECTION (LTBI) REPORT FORM SOUTH DAKOTA DEPARTMENT OF HEALTH

		PORTABLE TB RISK FACTO with latent TB infection wh		that apply) t one of the following risk factors:	
	Persons evaluated fo Immunosuppressive Radiographic evidence Children less than 5 y HIV infection Close contact (Defined	ce of prior TB	na therapy 12 months)	 □ Diabetes □ Renal dialysis □ Silicosis □ Organ transplant □ Head and neck cancers □ Leukemia □ Hodgkin's disease 	
	Report by fax: (60 TB Program questions: 1-8	05) 773-5509 800-592-1861 or	Report by mai	I: Tuberculosis Control Program South Dakota Department of Health 615 East 4 th Street	
-	(60	05) 773-3737		Pierre, SD 57501	
1.	PATIENT DEMOGRAPHIC	CS			
	Last Name	Firs Day	et Name te of Birth	Age	
	Home phone	Work phone		Cell phone	
	Parent/Guardian	Employer		Telephone #	
	<u>Gender</u> ☐ Male ☐ Female	Race Native American	☐ Black ☐ As	<u>Ethnicity</u> ian ☐ Non-Hispanic ☐ Hispanic	
	Foreign Born:	s If yes, country of birth			
	Clinic Name	Telephone # _		*(Required if foreign-born)	
	Physician	Fax #		Patient WeightLbs.	
_					
2.	. TB SCREENING INFORMATION				
	Date of test Date of bl			feron Gamma Release Assay) d collection ositive Negative Indeterminate	
	Classification: ☐ Reactor ☐ Conver ☐ Contact	tor Date of last negative test -		mm	
3.	CHEST X-RAY INFORMA	TION			
	Date of the chest X-ray	Results_			
1.	TREATMENT INFORMAT	ION			
	Starting Treatment?	□ No, Reason why		Yes, Date started	
	Therapy Prescribed:	□ INH and Rifapentine (3HP)		Once weekly for 12 doses (3 months)	
		□ Rifampin		Daily for 4 months	
		□ Isoniazid (INH)		□ Daily or □ Twice weekly □ 6 months or □ 9 months	
		□ INH and Rifampin (3HR)	[Daily for 3 months	
		□ Vitamin B-6	[□ Daily □ Weekly □ Twice weekly □ 3 months □ 6 months □ 9 months	_
	Medication Provider:				